

AD-A091 748

ROCHESTER UNIV NY DEPT OF CHEMISTRY

F/G 7/4

HIGH-ENERGY POSITRON IONIZATION OF ADSORBED SPECIES IN THE IMPU--ETC(U)

OCT 80 K LAM, T F GEORGE

N00014-80-C-0472

NL


UNCLASSIFIED

TR-7

1 of 1
2 - 100



END
DATE
FILMED
81-1-1
DTIC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

AD A091748

LEVEL

12

OFFICE OF NAVAL RESEARCH

Contract N00014-80-C-0472

Task No. NR 056-749

TECHNICAL REPORT NO. 7

High-Energy Positron Ionization of Adsorbed

Species in the Impulse Approximation

by

Kai-Shue Lam and Thomas F. George

in

The Journal of Physical Chemistry

University of Rochester
Department of Chemistry
Rochester, New York 14627

October, 1980

Reproduction in whole or in part is permitted for
any purpose of the United States Government

This document has been approved for public release
and sale; its distribution is unlimited

DTIC
ELECTED
NOV 18 1980

DDC FILE COPY

8011 10 144

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

14 TR-7

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 7	2. GOVT ACCESSION NO. AD-A091748	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) High-Energy Positron Ionization of Adsorbed Species in the Impulse Approximation		5. TYPE OF REPORT & PERIOD COVERED Interim Technical Report
6. AUTHOR(s) Kai-Shue Lam and Thomas F. George		7. PERFORMING ORG. REPORT NUMBER
8. PERFORMING ORGANIZATION NAME AND ADDRESS University of Rochester Department of Chemistry Rochester, New York 14627		9. CONTRACT OR GRANT NUMBER(s) N00014-80-C-0472
10. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Chemistry Program Code 472 Arlington, Virginia 22217		11. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 056-749
12. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		13. REPORT DATE October 1980
14. DISTRIBUTION STATEMENT (of this Report) This document has been approved for public release and sale; its distribution is unlimited.		15. NUMBER OF PAGES 11
16. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		17. SECURITY CLASS. (of this report) Unclassified
18. SUPPLEMENTARY NOTES Prepared for publication in The Journal of Physical Chemistry, in press (1981)		19. DECLASSIFICATION/DOWNGRADING SCHEDULE
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) POSITRON-SURFACE COLLISIONS THEORETICAL STUDY IONIZATION OF ADSPECIES IMPULSE APPROXIMATION HIGH ENERGY H/LI ADSPECIES/SURFACE SYSTEM		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) High-Energy Positron Ionization of adsorbed species is treated using the one-electron group-orbital linear-chain model and within the impulse approximation. Inter-orbital interference effects are found to be significant, leading to quenching (towards the low impact energy regime) and enhancement (at high impact energies) of ionization compared to positron gas-phase ionization.		

DD FORM 1473 JAN 73

EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102-LF-014-6601

Unclassified 307260
SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

J. Phys. Chem., in press.

High-Energy Positron Ionization of Adsorbed
Species in the Impulse Approximation

Kai-Shue Lam and Thomas F. George

Department of Chemistry
University of Rochester
Rochester, New York 14627

Abstract. High-Energy Positron Ionization of adsorbed species is treated using the one-electron group-orbital linear-chain model and within the impulse approximation. Inter-orbital interference effects are found to be significant, leading to quenching (towards the low impact energy regime) and enhancement (at high impact energies) of ionization compared to positron gas-phase ionization.

Accession For		<input checked="checked" type="checkbox"/>
NTIS GR&I		<input type="checkbox"/>
DTIC TAB		<input type="checkbox"/>
Unannounced		
Justification		
By		
Distribution/		
Availability Codes		
Avail and/or		
Special		
Dist		
HA		

I. Introduction

Recently, positron sources (available beam energy up to ~400 eV) have come to be of value in the diffraction studies of solid surfaces.¹ These studies complement the traditional low-energy electron diffraction (LEED) ones and point to new directions in the use of positrons as excitation or ionizing agents in the study of chemisorption bond-characters. Unlike electrons, positrons lead to no exchange effects during collision and thus allow for simplifications in theoretical analyses, provided the effects of positronium formation can be neglected (which should be a good approximation for beam energies of several hundred eV). Furthermore, in ionizing situations, the distinguishability of the product particles (positrons and electrons) also facilitates the measurement of ionization cross-sections.

Here we report on a theoretical treatment of high-energy positron ionization of adsorbed species based on the one-electron group-orbital linear-chain (OEGOLC) model of the adatom-surface system (with only nearest-neighbor interactions),^{2,3} and the impulse approximation (IA)⁴ for the description of the collision process. Indirect effects of the solid substrate on the ionization process, such as electron- and positron-phonon interactions, positron diffusion within the solid, positronium formation and positron annihilation, etc. will be ignored. Such effects are either expected to be weak compared to direct positron-electron interaction, or are negligible under the conditions of the high impact energies considered here (50-400 eV). Our main purpose is to investigate the effects of orbital mixing in the chemisorption bond on ionization. Hence we will focus on systems with

a localized adatom-surface (chemisorption) bond (corresponding to a localized state with energy lying outside a band). In this case, if the surface coverage is large enough, the ejected electrons will be mainly from near the surface, i.e., from the localized orbitals, and these electrons would contribute predominantly to the ionization cross-section compared to the "secondary" non-localized electrons. In our model, the positron is accordingly assumed to interact only with electrons in the localized bound state. The OEGOLC model, though somewhat qualitative, is the simplest one capable of producing a localized state. The IA, on the other hand, has been applied with considerable success to complex collision events such as high-energy gas-phase collisional ionization.⁵ Our calculations indicate that inter-orbital interference effects in the group orbital picture are significant in the ionization process: there is a slight quenching towards the low energy regime (< 90 eV) and considerable enhancement at high energies, compared to gas-phase ionization.

II. Theory

(a) The ionization cross-section

The physical process considered here is the collision between an incident positron (with beam energy considerably in excess of any bound-electron energies) and an adatom, and the subsequent single-ionization of the adatom. The IA assumes that, during collision, the interaction between the bound electrons of the adatom-surface system is "turned off". This implies that the incident positron can only spend a very short time in the field of the bound electrons. The electron interacting directly with the positron is "momentarily"

free and has a momentum (\vec{k}) distribution given by the Fourier transform $\tilde{\psi}_0(\vec{k})$ of the one-electron chemisorption wave function. The positron-electron system is then described by a positive-energy Coulomb wave function⁶ weighted by $\tilde{\psi}_0(\vec{k})$. If \vec{r}_1 is the positron coordinate, and \vec{r}_2 and \vec{r}_3 are coordinates of the electrons in the localized state, the positron-electron system wave function in the IA is thus given by (in what follows, all quantities will be expressed in terms of atomic units)

$$\begin{aligned} &\Psi(\vec{r}_1, \vec{r}_2, \vec{r}_3) \\ &= \frac{1}{\sqrt{2}} \{ \psi_0(\vec{r}_2) \int d^3k_3 \tilde{\psi}_0(\vec{k}_3) \psi_{\vec{k}_1, \vec{k}_3}(\vec{r}_1, \vec{r}_3) + \psi_0(\vec{r}_3) \int d^3k_2 \tilde{\psi}_0(\vec{k}_2) \psi_{\vec{k}_1, \vec{k}_2}(\vec{r}_1, \vec{r}_2) \}. \end{aligned} \quad (1)$$

In (1), \vec{k}_1 is the wave vector of the incident positron, $\psi_0(\vec{r})$ is the one-electron chemisorption wave function of the localized state, $\tilde{\psi}_0(\vec{k})$ is its Fourier transform, referred to above, and $\psi_{\vec{k}_i, \vec{k}_j}(\vec{r}_i, \vec{r}_j)$ is the Coulomb wave function for a positron-electron pair given by (on factoring into CM and relative coordinates)

$$\psi_{\vec{k}_1, \vec{k}_2}(\vec{r}_1, \vec{r}_2) = \frac{1}{(2\pi)^3} \exp\left\{\frac{1}{2}i(\vec{k}_1 + \vec{k}_2) \cdot (\vec{r}_1 + \vec{r}_2)\right\} \psi_{\frac{1}{2}(\vec{k}_1 - \vec{k}_2)}(\vec{r}_1 - \vec{r}_2) \quad (2)$$

where

$$\psi_{\vec{k}}(\vec{r}) = e^{\pi/2k} \Gamma\left(1 - \frac{i}{k}\right) e^{i\vec{k} \cdot \vec{r}} {}_1F_1\left(\frac{i}{k}; 1; ikr - i\vec{k} \cdot \vec{r}\right). \quad (3)$$

$\psi_{\vec{k}}(\vec{r})$ is the wave function in the continuous spectrum for an attractive Coulomb potential; ${}_1F_1$ is the confluent hypergeometric function.

The final states of the positron and ejected electron can be approximated by plane waves, and the final-state wave function for the positron-electron system can be written as

$$u_f(\vec{r}_1, \vec{r}_2, \vec{r}_3) = \frac{1}{\sqrt{2}} \frac{e^{i\vec{k}'_1 \cdot \vec{r}_1}}{(2\pi)^3} \{ e^{i\vec{k}'_2 \cdot \vec{r}_2} \psi_0(\vec{r}_3) + e^{i\vec{k}'_2 \cdot \vec{r}_3} \psi_0(\vec{r}_2) \}, \quad (4)$$

where \vec{k}'_1 and \vec{k}'_2 are the wave vectors for the scattered positron and emitted electron, respectively. The single-ionization scattering amplitude f is then given by

$$f = - \frac{1}{\sqrt{2\pi}} (T_2 + T_3), \quad (5)$$

where T_i , the transition matrix elements, can be expressed as

$$T_i = \int d^3r_1 d^3r_2 d^3r_3 u_f^*(\vec{r}_1, \vec{r}_2, \vec{r}_3) \left(-\frac{1}{r_{1i}}\right) \psi(\vec{r}_1, \vec{r}_2, \vec{r}_3); \quad (6)$$

and $r_{ij} = |\vec{r}_i - \vec{r}_j|$. Using (1) and (4) in (6) and expanding, we obtain a sum of eight terms for f , each of which represents different initial and final configurations for the positron-electron system coupled by a distinct $1/r_{ij}$ Coulomb force. Of these, in the Born approximation, four vanish and four are equal to one another, say, T . Based on this observation, we introduce the approximation that

$$T_2 + T_3 = 4T \quad (7)$$

where T is still calculated within the IA:

$$T = -\left(\frac{1}{2}\right) \frac{1}{(2\pi)^3} \int d^3r_1 d^3r_2 e^{-i\vec{k}'_1 \cdot \vec{r}_1 - i\vec{k}'_2 \cdot \vec{r}_2} \frac{1}{r_{12}} \int d^3k_2 \tilde{\psi}_0(\vec{k}_2) \psi_{\vec{k}_1, \vec{k}_2}(\vec{r}_1, \vec{r}_2). \quad (8)$$

Using (2) in (8), we obtain

$$T_2 + T_3 = -\frac{2}{(2\pi)^3} \tilde{\psi}_0(\vec{k}'_1 + \vec{k}'_2 - \vec{k}_1) \int d^3r \frac{e^{-i\left(\frac{\vec{k}'_1 - \vec{k}'_2}{2}\right) \cdot \vec{r}}}{r} \psi_{\frac{1}{2}(\vec{k}_1 - \vec{k}'_1 - \vec{k}'_2)}(\vec{r}). \quad (9)$$

The integral in (9) has been evaluated by several authors,⁷ and gives

$$T_2 + T_3 = -\frac{4}{(2\pi)^2} \tilde{\psi}_0(\vec{k}'_1 + \vec{k}'_2 - \vec{k}_1) \exp(\pi/|2\vec{k}_1 - \vec{k}'_1 - \vec{k}'_2|) \\ \times \frac{\Gamma(1-2i/|2\vec{k}_1 - \vec{k}'_1 - \vec{k}'_2|)}{(\vec{k}_1 - \vec{k}'_1)^2} \{ e^{-i\pi \frac{(\vec{k}_1 - \vec{k}'_1) \cdot (\vec{k}_1 - \vec{k}'_2)}{(\vec{k}_1 - \vec{k}'_1)^2}} \}^{-2i/|2\vec{k}_1 - \vec{k}'_1 - \vec{k}'_2|} \quad (10)$$

The wave vectors are required to satisfy the condition of energy conservation: $\kappa^2 = k_1^2 - 2|\epsilon| = k_1'^2 + k_2'^2$, where $-|\epsilon|$ is the bound energy of the localized electron. The total single-ionization cross section σ is then given by

$$\sigma = \frac{1}{k_1 k_2'} \int_{k_2' < \kappa} d^3 k_2' d\Omega_1 |f(\vec{k}_1', \vec{k}_2')|^2. \quad (11)$$

In (11), $d\Omega_1'$ designates the directions of the scattered positron. Only "backward-scattered" directions of \vec{k}_2' need be considered since these are the only observable ones. The scattering amplitude used in (11) is then given by

$$|f(\vec{k}_1', \vec{k}_2')|^2 = \frac{2^4 (2\pi)^2 |\tilde{\psi}_0(\vec{k}_1' + \vec{k}_2' - \vec{k}_1)|^2 \exp(-2\pi/|2\vec{k}_1 - \vec{k}_1' - \vec{k}_2'|)}{|2\vec{k}_1 - \vec{k}_1' - \vec{k}_2'| |\vec{k}_1 - \vec{k}_1'|^4 \sinh(2\pi/|2\vec{k}_1 - \vec{k}_1' - \vec{k}_2'|)} \quad (12)$$

(b) The chemisorption wave function

The chemisorption wave function in the OEGOLC model can be written, using the LCAO approximation, as

$$\psi_0(\vec{r}) = C_\lambda \phi_a(\vec{r}) + \sum_{m=0}^{N-1} C_m \phi(\vec{r} + (\lambda+m)\hat{a}\hat{z}), \quad (13)$$

where \hat{z} is in the direction of the chain (outward from the surface), a is the inter-atomic distance in the linear chain, λa is the distance between the adatom and the first substrate atom, $N+1$ is the total number of substrate atoms in the chain, and \vec{r} is measured from the nucleus of the adatom. We assume that only two types of orbitals, ϕ_a (corresponding to the free adatom) and ϕ (corresponding to individual substrate atoms) enter into ψ_0 . [This assumption should be reasonable for most alkalis and some transition metals.] The coefficients C_λ and C_m depend on the following parameters:

ϵ_a (bound energy corresponding to ϕ_a), α (bound energy corresponding to ϕ), β' (interaction between the adatom and the first substrate atom) and β (nearest-neighbor interaction between the substrate atoms).

When (13) is used in the Schrödinger equation with a one-electron Hamiltonian, the vanishing of the secular determinant for the coefficients C_λ and C_m leads to the following result. A localized state (ψ_0 damped in the crystal) exists if the equation

$$(\cos\theta + \sin\theta \cot N\theta) \left(\frac{\alpha - \epsilon_a}{\beta} + 2\cos\theta \right) = \left(\frac{\beta'}{\beta} \right)^2 \quad (14)$$

has at least one complex solution for θ , ($\theta = i\xi$ or $\pi + i\xi$, $\xi > 0$ and real), with energy given by $\epsilon = \alpha \pm 2\beta \cosh\xi$. $|\tilde{\psi}_0|^2$ in (12) is then given by

$$\begin{aligned} |\tilde{\psi}_0(\vec{k})|^2 = & C_\lambda^2 \tilde{\phi}_a^2 + \tilde{\phi}^2 [1 - C_\lambda^2 + 2 \sum_{m' > m} C_{m'} C_m \cos\{a \vec{k} \cdot \hat{z}(m' - m)\}] \\ & + 2C_\lambda \tilde{\phi}_a \tilde{\phi} \sum_m C_m \cos\{(\lambda + m)a \vec{k} \cdot \hat{z}\}, \end{aligned} \quad (15)$$

where $\tilde{\phi}_a$ and $\tilde{\phi}$ are the Fourier transforms of ϕ_a and ϕ , respectively, and both are taken to be real. The secular equations for C_λ and C_m also lead to

$$C_\lambda = \left\{ \frac{\beta'/\beta}{(\alpha - \epsilon_a)/\beta \pm 2 \cosh \xi} \right\} C_0 \quad (16a)$$

and

$$C_m = \frac{\sinh\{(N-m)\xi\}}{\sinh N\xi} C_0 \quad (16b)$$

For a localized (damped wave function, the series in (15) terminates quickly. (15) displays clearly the inter-orbital interference effects.

III. Results and Discussion

Calculations have been carried out for the H(adatom)/Li(substrate) system with normally incident positrons. β and β' are estimated by the Wolfsberg-Helmholz formula,⁸ and the overlap integrals required therein are obtained from tabulations from Mulliken et al.⁹ ϕ_a and ϕ are taken to be the H 1s and Li 2s Slater AO's, respectively. The following parameters are then obtained (in a.u.): $\beta = -0.17$, $\beta' = -0.35$, $\epsilon = -0.76$. Table 1 gives the normalized coefficients C_λ and C_m for $N = 100$ computed using (16). To the accuracy stated, these results are identical to those for $N = 14$. The energy gap between the localized state and the bottom edge of the 2s band is 0.22 a.u.

The integrations in (11) are carried out numerically using the 6- and 16-point Gaussian quadratures, and the ionization cross sections are compared with those for the positron ionization of the H atom, also computed using the IA (Fig. 1). In the latter case, only a simple orbital enters into (12), and inter-orbital interference effects are absent. [For purposes of comparison, σ for the H atom case only include "backward" ionization directions.] Our results indicate that at high impact energies there is a distinct enhancement in ionization due to the interference effects of the group orbital, while at low energies these effects tend to quench the ionization. The peak cross section, however, occurs at a higher energy for the adatom case than for the gas-phase case. Even though the low energy results are less reliable due to the doubtful validity of the IA in this regime, the increasing probability of positronium formation, and hence positron annihilation,

also tend to decrease the ionization cross section. Hence the overall trend is expected to be quenching in the low-energy regime and enhancement in the high energy regime. The enhancement may be attributed to the fact that, at high energies, the positron probes deeper into the crystal.

TABLE 1. Normalized coefficients for the localized H/Li chemisorption wavefunction with bound energy $\epsilon = -0.76$ a.u.


m	$m=\lambda$	0	1	2	3	4	5	6	7	8
C_m	0.8005	0.5637	0.1917	0.0648	0.0220	0.0073	0.0025	0.0009	0.0003	0

Acknowledgments

This work was supported in part by the Office of Naval Research, and by the Air Force Office of Scientific Research (AFSC), United States Air Force, under Contract No. F49620-78-C-0005. The United States Government is authorized to reproduce and distribute reprints for governmental purposes notwithstanding any copyright notation hereon. One of us (TFG) acknowledges the Alfred P. Sloan Foundation for a Research Fellowship and the Camille and Henry Dreyfus Foundation for a Teacher-Scholar Grant.

1. I. J. Rosenberg, A. H. Weiss, and K. F. Canter, Phys. Rev. Lett. 44, 1139 (1980), and references cited therein.
2. T. B. Grimley, Proc. Phys. Soc. (London) 72, 103 (1958).
3. J. P. Muscat and D. M. Newns, Prog. Surf. Sci. 9, 1 (1978).
4. For an application of the IA to gas-phase electron-atom scattering, see, for example, R. Akerib and S. Borowitz, Phys. Rev. 122, 1177 (1961).
5. H. Nakamura, T. Shirai, and Y. Nakai, Phys. Rev. A 17, 1892 (1978).
6. See, for example, N. F. Mott and H. S. W. Massey, "The Theory of Atomic Collisions," 3rd ed. (Oxford, London, 1965), p. 56.
7. See, for example, A. Nordsieck, Phys. Rev. 93, 785 (1954).
8. M. Wolfsberg and L. Helmholz, J. Chem. Phys. 20, 837 (1952).
9. R. S. Mulliken, C. A. Rieke, D. Orloff, and H. Orloff, J. Chem. Phys. 17, 1248 (1949).

Fig. 1. Positron single-ionization cross sections (σ) for H/Li and H atom as a function of incident positron energy (E). $a_0 = 1$ Bohr. —, H/Li; ----, H atom. In both cases only "backward" ionization directions are included.



TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 472 800 North Quincy Street Arlington, Virginia 22217	2	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, N.C. 27709	1
ONR Branch Office Attn: Dr. George Sandoz 536 S. Clark Street Chicago, Illinois 60605	1	Naval Ocean Systems Center Attn: Mr. Joe McCartney San Diego, California 92152	1
ONR Area Office Attn: Scientific Dept. 715 Broadway New York, New York 10003	1	Naval Weapons Center Attn: Dr. A. B. Amster, Chemistry Division China Lake, California 93555	1
ONR Western Regional Office 1030 East Green Street Pasadena, California 91106	1	Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko Port Hueneme, California 93401	1
ONR Eastern/Central Regional Office Attn: Dr. L. H. Peebles Building 114, Section D 666 Summer Street Boston, Massachusetts 02210	1	Department of Physics & Chemistry Naval Postgraduate School Monterey, California 93940	1
Director, Naval Research Laboratory Attn: Code 6100 Washington, D.C. 20390	1	Dr. A. L. Slafkosky Scientific Advisor Commandant of the Marine Corps (Code RD-1) Washington, D.C. 20380	1
The Assistant Secretary of the Navy (RE&S) Department of the Navy Room 4E736, Pentagon Washington, D.C. 20350	1	Office of Naval Research Attn: Dr. Richard S. Miller 800 N. Quincy Street Arlington, Virginia 22217	1
Commander, Naval Air Systems Command Attn: Code 310C (H. Rosenwasser) Department of the Navy Washington, D.C. 20360	1	Naval Ship Research and Development Center Attn: Dr. G. Bosmajian, Applied Chemistry Division Annapolis, Maryland 21401	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12	Naval Ocean Systems Center Attn: Dr. S. Yamamoto, Marine Sciences Division San Diego, California 91232	1
Dr. Fred Saalfeld Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1

TECHNICAL REPORT DISTRIBUTION LIST, GENNo.
Copies

Dr. Rudolph J. Marcus
Office of Naval Research
Scientific Liaison Group
American Embassy
APO San Francisco 96503

1

Mr. James Kelley
DTNSRDC Code 2803
Annapolis, Maryland 21402

1

Dr. David L. Nelson
Chemistry Program
Office of Naval Research
800 North Quincy Street
Arlington, Virginia 22217

1

TECHNICAL REPORT DISTRIBUTION LIST, 056

	<u>No. Copies</u>		<u>No. Copies</u>
Dr. D. A. Vroom IRT P.O. Box 80817 San Diego, California 92138	1	Dr. C. P. Flynn Department of Physics University of Illinois Urbana, Illinois 61801	1
Dr. G. A. Somorjai Department of Chemistry University of California Berkeley, California 94720	1	Dr. W. Kohn Department of Physics University of California (San Diego) LaJolla, California 92037	1
Dr. L. N. Jarvis Surface Chemistry Division 4555 Overlook Avenue, S.W. Washington, D.C. 20375	1	Dr. R. L. Park Director, Center of Materials Research University of Maryland College Park, Maryland 20742	1
Dr. J. B. Hudson Materials Division Rensselaer Polytechnic Institute Troy, New York 12181	1	Dr. W. T. Peria Electrical Engineering Department University of Minnesota Minneapolis, Minnesota 55455	1
Dr. John T. Yates Surface Chemistry Section National Bureau of Standards Department of Commerce Washington, D.C. 20234	1	Dr. Narkis Tzoar City University of New York Convent Avenue at 138th Street New York, New York 10031	1
Dr. Theodore E. Madey Surface Chemistry Section Department of Commerce National Bureau of Standards Washington, D.C. 20234	1	Dr. Chia-wei Woo Department of Physics Northwestern University Evanston, Illinois 60201	1
Dr. J. M. White Department of Chemistry University of Texas Austin, Texas 78712	1	Dr. D. C. Mattis Polytechnic Institute of New York 333 Jay Street Brooklyn, New York 11201	1
Dr. Keith H. Johnson Department of Metallurgy and Materials Science Massachusetts Institute of Technology Cambridge, Massachusetts 02139	1	Dr. Robert M. Hexter Department of Chemistry University of Minnesota Minneapolis, Minnesota 55455	1
Dr. J. E. Demuth IBM Corporation Thomas J. Watson Research Center P.O. Box 218 Yorktown Heights, New York 10598	1	Dr. R. P. Van Duyne Chemistry Department Northwestern University Evanston, Illinois 60201	1

TECHNICAL REPORT DISTRIBUTION LIST, 056

	<u>No.</u> <u>Copies</u>		<u>No.</u> <u>Copies</u>
Dr. M. G. Lagally Department of Metallurgical and Mining Engineering University of Wisconsin Madison, Wisconsin 53706	1	Dr. J. Osteryoung Chemistry Department SUNY, Buffalo Buffalo, New York 14214	1
Dr. Robert Gomer Department of Chemistry James Franck Institute 5640 Ellis Avenue Chicago, Illinois 60637	1	Dr. G. Rubloff I.B.M. Thomas J. Watson Research Center P. O. Box 218 Yorktown Heights, New York 10598	1
Dr. R. G. Wallis Department of Physics University of California, Irvine Irvine, California 92664	1	Dr. J. A. Gardner Department of Physics Oregon State University Corvallis, Oregon 97331	1
Dr. D. Ramaker Chemistry Department George Washington University Washington, D.C. 20052	1	Dr. G. D. Stein Mechanical Engineering Department Northwestern University Evanston, Illinois 60201	1
Dr. P. Hansma Chemistry Department University of California, Santa Barbara Santa Barbara, California 93106	1	Dr. K. G. Spears Chemistry Department Northwestern University Evanston, Illinois 60201	1
Dr. P. Hendra Chemistry Department Southampton University England SO9 5NH	1	Dr. R. W. Plummer University of Pennsylvania Department of Physics Philadelphia, Pennsylvania 19104	1
Professor P. Skell Chemistry Department Pennsylvania State University University Park, Pennsylvania 16802	1	Dr. E. Yeager Department of Chemistry Case Western Reserve University Cleveland, Ohio 41106	2
Dr. J. C. Hemminger Chemistry Department University of California, Irvine Irvine, California 92717	1	Professor George H. Morrison Cornell University Department of Chemistry Ithaca, New York 14853	1
Dr. Martin Fleischmann Department of Chemistry Southampton University Southampton SO9 5NH Hampshire, England	1	Professor N. Winograd Pennsylvania State University Chemistry Department University Park, Pennsylvania 16802	1
		Professor Thomas F. George Department of Chemistry University of Rochester Rochester, New York 14627	1

